

Accepted Manuscript

Title: Biomarkers associated with obstructive sleep apnea and morbidities: a scoping review

Author: Graziela De Luca Canto, Camila Pachêco-Pereira, Secil Aydinoz, Paul W. Major, Carlos Flores-Mir, David Gozal

PII: S1389-9457(15)00029-5
DOI: <http://dx.doi.org/doi: 10.1016/j.sleep.2014.12.007>
Reference: SLEEP 2633

To appear in: *Sleep Medicine*

Received date: 10-10-2014
Revised date: 16-12-2014
Accepted date: 27-12-2014

Please cite this article as: Graziela De Luca Canto, Camila Pachêco-Pereira, Secil Aydinoz, Paul W. Major, Carlos Flores-Mir, David Gozal, Biomarkers associated with obstructive sleep apnea and morbidities: a scoping review, *Sleep Medicine* (2015), <http://dx.doi.org/doi: 10.1016/j.sleep.2014.12.007>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Biomarkers associated with obstructive sleep apnea and morbidities: a scoping review

Authors

Graziela De Luca Canto; Camila Pachêco-Pereira; Secil Aydinöz; Paul W. Major; Carlos Flores-Mir; David Gozal

Dr. De Luca Canto is adjunct professor, Department of Dentistry, Federal University of Santa Catarina, Florianópolis, SC, Brazil and Adjunct Assistant Professor at School of Dentistry, Faculty of Medicine and Dentistry, University of Alberta, Edmonton Clinic Health Academy, 11405-87 Avenue, Edmonton, Alberta, Canada T6G 1C9. Email: graziela.canto@ufsc.br

Dr. Pachêco-Pereira is Masters student, School of Dentistry, Faculty of Medicine and Dentistry, School of Dentistry, University of Alberta, Edmonton Clinic Health Academy, 11405-87 Avenue, Edmonton, Alberta, Canada T6G 1C9. Email: cppereir@ualberta.ca

Dr. Aydinöz is Associate Professor of Pediatrics at GATA Haydarpasa Teaching Hospital, Istanbul, Turkey, and Visiting Scholar in the Section of Pediatric Sleep Medicine, Department of Pediatrics, The University of Chicago, USA. Email: saydinöz@gmail.com

Dr. Major is Professor and Chair, School of Dentistry, Senior Associate Dean, Dental Affairs, School of Dentistry, Faculty of Medicine and Dentistry, University of Alberta. Edmonton Clinic Health Academy, 11405-87 Avenue, Edmonton, Alberta, Canada T6G 1C9. Email: major@ualberta.ca

Dr. Flores-Mir is Associate Professor, Division Head of Orthodontics, School of Dentistry, Faculty of Medicine and Dentistry, University of Alberta, Edmonton Clinic Health Academy, 11405-87 Avenue, Edmonton, Alberta, Canada T6G 1C9. Email: cf1@ualberta.ca

Dr. David Gozal is Professor, Department of Pediatrics, Pritzker School of Medicine, Biological Sciences Division, The University of Chicago, 5721 S. Maryland Avenue, MC 8000, Suite K-160 Chicago, IL 60637, USA.

Email: dgozal@uchicago.edu

Conflict of interest: The authors have no conflict of interest to declare.

Financial support: DG is supported by NIH grant HL65270-12.

Corresponding author:

David Gozal, MD

Department of Pediatrics

The University of Chicago

5721 S. Maryland Avenue, MC 8000, Suite K-160

Chicago, IL 60637

(773) 702-3360

(773) 926-0756- FAX

Email: dgozal@uchicago.edu

Highlights

- The majority of OSA-related biomarkers have relied on blood-based assays.
- Plasma IL-6 and CRP emerge as promising morbidity biomarkers in both adults and children.
- Urinary neurotransmitters identify cognitive morbidity in children with OSA.

Abstract

Objective: To map potential biomarkers of obstructive sleep apnea (OSA)-associated morbidities in both adults and children, to identify gaps in current evidence, and to determine the value of conducting a full systematic review.

Methods: A scoping review was undertaken of studies in patients with OSA that evaluated the potential value of biological markers in identifying OSA-associated morbidities. Retained articles were only those studies whose main objective was to identify morbidity biomarkers in subjects with OSA, the latter being confirmed with a full overnight polysomnography (PSG) in a laboratory or at-home settings. The methodology of the selected studies was classified using an adaptation of the evidence quality criteria recommended by the American Academy of Pediatrics. Additionally the biomarkers were categorized according to their potential clinical applicability.

Results: 572 citations were identified of which 48 met inclusion criteria. Thirty-four studies were conducted in adults and 14 involved children. Most of the studies evaluated blood biomarkers, and presented 31 potential diagnostic biomarkers.

Conclusion: The majority of studies were performed explored blood-based biomarkers, with most not identifying definitive morbidity biomarkers. Of the potentially promising morbidity biomarkers, plasma IL-6 and high sensitivity C-reactive protein appear to exhibit a favorable

profile, and may discriminate OSA patients with and without morbidities in both adults, as well as MRP 8/14 in children. Urinary neurotransmitters may also provide a good tool for screening OSA cognitive morbidity in children.

Keywords: review, sleep apnea, biomarkers, morbidity

Abbreviations

OSA=obstructive sleep apnea

CVD=cardiovascular disease

DTA= diagnostic test accuracy

EDS= excessive daytime sleepiness

PSG= polysomnography

EBC= exhaled breath condensate

RR=risk ratio

OR=odds ratio

AHI=apnea hypopnea index

OAI= obstructive apnea index

RDI= respiratory disturbance index

Hr/TST=hour of total sleep time

ROC= receiver operating characteristic

AASM=American Academy of Sleep Medicine

hsCRP=high-sensitivity C-reactive protein

IGF= Insulin-like growth factor

MRP=myeloid-related protein

IL-6=interleukin-6

1 Introduction

Obstructive sleep apnea (OSA) has now been recognized as a major public health issue with potential society-wide consequences involving car or work-related accidents, cognitive and behavioral deficits impairing work performance, and potentially leading to cardiovascular and metabolic dysfunction.¹ Indeed, OSA has been associated with serious morbidities such as endothelial dysfunction^{2,3}, hypertension⁴, cardiovascular disease (CVD)⁵⁻⁷, cognitive and behavioral dysfunction⁸, metabolic disorders such as insulin resistance⁹, diabetes¹⁰, and dyslipidemias¹¹, erectile dysfunction in men,¹² nocturnal enuresis in children¹³, and excessive daytime sleepiness (EDS)^{14,15}. Consequently, healthcare costs are substantially increased in patients with OSA, accounting either directly or via its associated morbidities for a substantial proportion of all medical-related costs.¹⁶⁻¹⁸

In adults, the prevalence of OSA varies widely, from 14.7% to 36.5%, depending on age, gender and ethnicity.¹⁹ It is generally higher in males,¹⁹ and although the prevalence in Hispanics is similar to white Caucasians, it is significantly higher in African American or Asians.¹⁹ In children, the prevalence of OSA is reported as ranging between 1 and 4%.^{20,21}

Current diagnostic approaches range from exclusively using clinical presentation and physical

examination to the current ‘gold standard’, the overnight polysomnography (PSG). However, the measures derived from PSG are poor predictors of OSA-associated morbidities.²² In other words, two patients with similar OSA severity may present with markedly different clinical phenotypes, whereby one will manifest substantial end-organ morbidities related to the presence of OSA, while all such features are absent in the other. The phenotypic variance in the clinical morbidity of OSA has therefore prompted exploration of biomarkers that would enable the identification of the more “vulnerable” patients, who would more likely benefit from timely and targeted therapeutic interventions. In other words, such studies explored the opportunity to enable incorporation of morbidity biomarkers into well-defined and validated clinical algorithms.²² The search for appropriate biomarkers becomes therefore critical. The discovery of an ideal biomarker for OSA-associated morbidity has the potential to provide information related to prognosis and response to treatment.⁵ Ideal biomarkers should be highly sensitive and specific for OSA-induced end-organ dysfunction, should be involved in an important causal pathway, so that changes in the biomarker levels in the context of OSA treatment reliably predict improvements in the specific end-organ outcome.²³

Several different morbidity biomarkers have been proposed for OSA over the last 12 years. However, to the best of our knowledge, no scoping review has thus far been conducted to understand what we know about the use of biomarkers in the identification and management of OSA-associated morbidities. Therefore, the purpose of this scoping review was to map our current understanding regarding any of the putative biomarkers that have been thus far investigated regarding their potential association or predictive ability of OSA-associated morbidities in both adults and children, to identify gaps in the research, and to determine the value of conducting a full systematic review related to this topic.

Accepted Manuscript

2 Methods

This scoping review was performed while adhering to Arksey and O'Malley's scoping review proposed reporting framework.²⁴

2.1 Research question

A scoping review of studies in subjects with OSA that evaluated the potential diagnostic value of biological markers (blood, exhaled breath condensate (EBC), salivary, and urinary) in the identification of those patients with morbidities associated with the underlying disease was undertaken.

2.2 Identifying relevant studies

Inclusion criteria

Retained articles were only those studies which objective was to identify morbidity-related biomarkers in patients with OSA (cognitive, excessive sleepiness, cardiovascular, and/or metabolic), with the diagnosis being confirmed with a full overnight polysomnography (PSG) in either the laboratory or home setting. Only studies that performed PSG in all subjects were included. Studies that assessed the impact of treatment were also included. Lack of a control group was accepted. Only studies in English, Spanish and Portuguese were considered.

Exclusion criteria

Studies using daytime PSG or respiratory polygraphy were not considered. Studies in which the cohort included syndromic patients (e.g., Down syndrome, craniofacial anomalies, neuromuscular disorders, etc.) or patients with a primary disease for which OSA prevalence is being investigated (e.g., patients with kidney disease, and/or rheumatologic conditions), were discarded. Reviews, letters, conference abstracts and personal opinions were not considered. Studies about central apnea were not included.

In phase 2, we excluded studies that explored OSA only, but did not assess OSA-associated morbidities.

Detailed individual search strategies for each of the following bibliographic databases were developed: Cochrane, EMBASE, MEDLINE, PubMed, and LILACS. A partial grey literature search was undertaken using Google Scholar. The end search date for all database searches was March 20, 2014. The references cited in the selected articles were also checked for any references that could have been missed during the electronic database searches. Additional studies that were already known by the authors but were not identified in the searches were also included.

Appropriate truncation and word combinations were selected and were adapted for each database search. (Appendix A) All references were managed by reference manager software (RefWorks – COS, a business unit of ProQuest, LLC. ©7200 Wisconsin Avenue, Suite 601 Bethesda, MD 20866 USA) and duplicate hits were removed.

2.3 Study Selection

The selection was completed in two phases. In phase 1, two reviewers independently reviewed the titles and abstracts of all identified electronic database citations (GDL and CPP). A third author was involved when required to make a final decision (SA). Any studies that appeared not to fulfill the inclusion criteria were discarded. In phase 2, the same selection criteria were applied to the full articles to confirm their eligibility. The same two reviewers (GDL and CPP) independently participated in phase 2. The reference list of included articles was revised by one examiner (GDL). The articles selected were read by both examiners (GDL and CPP). Any disagreement in either phase was resolved by discussion and mutual agreement between the three reviewers (GDL, CPP, SA). A fourth author, expert in sleep medicine (DG) was involved when controversy arose before making a final decision. Final selection was always based on the full-

text of the publication.

2.4 Charting the data

For the included studies the following information was recorded: year of publication, author, country, sample characteristics, name and type of OSA-related morbidity biomarkers, OSA diagnostic cut-off value at PSG, results and main conclusion. Authors of potentially eligible full-articles were contacted as necessary to provide further details about their studies.

One author (GDL) collected the required information from the selected articles. A second author (CPP) crosschecked all the collected information. Again, any disagreement in either phase was resolved by discussion and mutual agreement between the three reviewers (GDL, CPP, SA). A fourth author (DG) was involved, when required, to reach the final decision.

2.5 Level of Evidence

The methodology of selected studies was classified using a non-validated adaptation of the evidence quality criteria from American Academy of Pediatrics ²⁵. Two reviewers (GDL and CPP) independently classified the studies as A (well-designed prognostic or diagnostic studies on relevant population), B (prognostic or diagnostic studies with minor limitations, overwhelmingly consistent evidence from observational studies), C (observational studies (case-control and cohort design)). Disagreements were solved by a third reviewer (DG).

Additionally, the biomarker clinical application was classified as (1) potential clinical biomarker(s) of OSA morbidity; (2) inconclusive for clinical biomarker of OSA morbidity, and (3) evidence not supportive as potential biomarker(s). Two reviewers (GDL and CPP) independently classified the clinical application of biomarkers. A third reviewer (SA) revised the classification. Disagreements were further resolved by a fourth reviewer (DG).

2.6 Collating, summarizing and report the results

Any outcome measurement was considered (risk ratio (RR), odds ratio (OR) or risk difference for dichotomous outcomes; mean difference or standardized mean difference for continuous outcomes; sensitivity and specificity in diagnostic studies).

Accepted Manuscript

3 Results

3.1 Study Selection

During the initial phase, 572 citations were identified across the five electronic databases. After removing the duplicates articles, only 279 remained. A comprehensive evaluation of the abstracts resulted in the retention of 183 articles after phase 1. In addition 40 citations were identified in Google Scholar; however, only 4 of those citations met phase 1 inclusion criteria. One additional study was identified from the hand-search of reference lists of these studies. Six more studies were added from a content expert. Therefore, the final phase I count was 194 articles, including the 6 referred by an expert. After a full-text review, 146 studies were excluded. Reasons for exclusion can be found in Appendix B. At the end of the selection process only 48 articles were retained. A flow chart of the process of identification, inclusion, and exclusion of studies is shown in Figure 1.

Add Figure 1 here

3.2 Study Characteristics

The selected studies were grouped into two categories: children and adults. Thirty-four studies^{3,4,6,7,10,15,26-53} were conducted in adults and 14 in children^{2,8,9,13,14,54-62}.

The pediatric studies were published between 2007 and 2013 They were conducted in United States^{2,8,9,13,14,58-62, 56,57}, China⁵⁵ and Egypt⁵⁴. The OSA diagnostic criterion was established by AHI, OAI or RDI. When the authors used AHI, the diagnostic AHI ranged from AHI>1 to AHI >2 per hour of total sleep time (/hrTST). The majority of these studies assessed blood biomarkers, except for one study that studied blood and urinary biomarkers⁹, one that studied urinary biomarkers⁵⁹, and one EBC⁵⁴. In children, studies primarily focused cardiac diastolic dysfunction, insulin resistance, cognitive functional morbidity, enuresis, and vascular dysfunction. A summary of these studies' descriptive characteristics can be found in Table 1 and

Appendix C.

The studies in adults were published between 2002 and 2014. The majority was published in the United States^{26,28,45,46,53}, Turkey^{6,29,30,43}, and Japan^{42,44}. The OSA diagnostic criterion was established by AHI or RDI. When the authors used AHI, the AHI ranged from AHI >5 to AHI ≥ 30 /hrTST. Most of these reports studied blood biomarkers, three studies used both blood and urine^{3,48,53}, one used blood and saliva⁷, and one EBC⁴¹. Major morbidities explored were CVD and cerebrovascular injury. A summary of these studies' descriptive characteristics can be found in Table 2 and Appendix D.

Add Table 1 and Table 2 here

3.3 Level of Evidence

Almost all pediatrics studies were classified as B (prognostic or diagnostic studies with minor limitations, overwhelmingly consistent evidence from observational studies). Only one study was classified as C (case-control and cohort design)¹⁴ and two studies were classified as A (well designed prognostic or diagnostic studies on relevant population)^{8,61}. None of the studies however performed ROC analysis for their biomarkers of interest, except Kheirandish-Gozal et al⁵⁹ who explored combinatorial biomarker approaches using specific cut-offs for levels of urinary neurotransmitters and their overnight changes.

In adult studies, 29 studies were classified as B^{3,4,6,7,10,15,26-53}. Five studies were classified as C^{4,28,38(Kato, 2011 #82,44,50)}. None of the studies was classified as A.

Synthesis of Results

Thirty-one potential biomarkers were identified. Only two (hsCRP and IL-6) were evaluated in both adults and children (Table 3). Only one study⁵⁹ reported diagnostic test accuracy (DTA) measurements about the biomarkers evaluated. Kheirandish-Gozal⁵⁹ performed a neurocognitive battery assessing general cognitive ability (GCA) to explore overnight changes in

neurotransmitters in the urine in children with and without OSA. They concluded that pediatric OSA is associated with overnight increases in urinary concentrations of catecholamines and that combinatorial approaches using defined cutoffs in overnight changes in concentration of selected neurotransmitters in urine may predict both OSA and the presence of cognitive deficits.

Among the 17 countries that have been published about this topic, most of the studies originated from the USA (Table 4).

Add Tables 3 and 4 here

4 Discussion

The present scoping review investigated the available evidence aiming to identify biomarkers in subjects with OSA-associated morbidities.

OSA is one of the important independent risk factors for CVD⁵⁻⁷, including hypertension⁴, cerebrovascular disease³³, and endothelial dysfunction^{2,3} with diabetes⁶³, cognitive dysfunction⁸, excessive daytime sleepiness (EDS)^{14,15} being also investigated, albeit less frequently. The risk of mortality was pronounced particularly in those with moderate-to-severe OSA. The existence of OSA associated with morbidities provides the need to develop different and likely combined therapeutic approaches according to the patient phenotypic complexity. The close association between OSA and comorbidities may magnify the cardiovascular risk factors, aggravating mortality.⁶⁴

The overnight PSG has been considered as the actual gold standard for OSA diagnosis. However, one of its major limitations resides in its inability to accurately identify those patients with OSA who manifest end-organ morbidity as a consequence of the underlying OSA. A method allowing for the screening of OSA patients in a large scale, and enabling an accurate identification of those patients with specific morbidities could potentially revolutionize the field.⁶⁵ This urgent need to find an ideal biomarker for OSA specific morbidities could explain the large amount of studies that have somewhat addressed this topic since 2000. The ideal biomarker should have some characteristics, such as morbidity specificity, presence in all patients with the end-organ dysfunction of interest, reversibility following proper treatment, and detectability before patients develop several clinical manifestations. It should also reflect the severity of the clinical morbid manifestation, and provide an indicator of cumulative injury over time, as well as exhibit minimal overlap between no morbidity and morbidity.⁶⁶

Different groups of researchers have tried to identify morbidity-related biomarkers in children and adults with OSA throughout the world. The United States leads in such efforts in both adults and children. Interestingly, in some countries such as Turkey and Japan, studies have been performed studied in adults, but not in children.

It is also noteworthy the wide range variation in OSA diagnostic criteria. AHI is the most frequently used marker of OSA severity. Nevertheless, the use of AHI is associated with two major limitations. Firstly, the clinically valid cut-off for normal AHI is unclear in children. Secondly, no consensus has been achieved as to whether children with AHI values between the normal cut-off ($<1/\text{hrTST}$) and $5/\text{hrTST}$ should undergo adenotonsillectomy.⁶⁶ It is therefore difficult to define OSA when exclusively using PSG-based criteria, a problem that has led to several experts and even clinical practitioners to state that “you will know that OSA is present when you see it”. This lack of consensus in the diagnostic criteria is the result of the relative dichotomy between PSG-derived measures and clinical symptoms. Children who are very symptomatic may present a “normal PSG” in the presence of habitual snoring. Conversely, asymptomatic snoring children may have concurrent and severe respiratory disturbance in their PSG.⁶⁷ To overcome some of this debate, the American Academy of Sleep Medicine (AASM)⁶⁸ has set a threshold of 15 events per hour with or without symptoms or 5 events per hour with symptoms for the diagnosis of OSA in adults⁶⁹

Interestingly, the majority of studies evaluated blood biomarkers. Few studies evaluated saliva, urine and/or EBC, a surprising finding considering that these approaches are noninvasive and are easily collected, especially in children.

When analyzing the level of evidence, only two studies were classified as A.^{8,61} To assess the magnitude of the systemic inflammatory response, as measured by high-sensitivity C-reactive

protein (hsCRP) serum levels, that may identify children with OSA at higher risk for cognitive morbidity, Gozal et al 2007⁸ recruited habitually snoring children and non-snoring children from the community. All children underwent PSG and neurocognitive testing and a fasting blood draw the next morning. Snoring children were divided into OSA and non-OSA groups, and children with OSA were further subdivided into those with two or more abnormal cognitive subtests and those with normal cognitive scores. Gozal and colleagues⁸ concluded that hsCRP levels are higher in children with OSA, and particularly in those who develop neurocognitive deficits, suggesting that the magnitude of the inflammatory responses elicited by OSA is a major determinant of increased risk for neurocognitive dysfunction. Indeed, among the 103 snoring children without OSA, hsCRP levels were 0.19 ± 0.07 mg/dl (p value not significant vs. control children). However, 12 of the snoring children without OSA exhibited cognitive deficits and their hsCRP levels were 0.33 ± 0.10 mg/dl ($p < 0.04$ vs. habitual snorers with normal cognitive scores). Among the 102 children with OSA, 57 had evidence of abnormal cognitive function, and these children were those whose hsCRP levels, namely 0.48 ± 0.12 mg/dl, markedly differed from not only control children ($p < 0.0001$) or snoring children ($p < 0.0001$), but also from the OSA children with preserved cognitive abilities (0.21 ± 0.08 mg/dl; $p < 0.002$).⁸ Gozal⁶¹ recruited children from the community and underwent overnight PSG to test if the lower IGF-1 serum levels in children with OSA would be associated with more severe neurocognitive morbidity compared with matched controls. They concluded that IGF-1 levels are higher in OSA children, mainly in those who do not manifest neurocognitive deficits.

Potential biomarkers in children have mainly focused on the investigation of myeloid-related protein (MRP) 8/14, and urinary neurotransmitters. On the other hand, studies in adults with OSA have focused on hsCRP and IL-6, and the latter 2 candidates were the only ones evaluated in both

adults and children. We should emphasize that only one study presented DTA measurements, even if it is now widely recognized that the lack of DTA results is concerning and should not be viewed as acceptable in biomarker studies.⁷⁰ However, the findings reported herein could provide the basis for future multicenter research efforts. In addition, there is a critical need for future studies to assess and provide a cut-off value for each of the biomarkers being evaluated and thus enable comparisons across studies as well as across age groups (e.g., adults vs. children). Notwithstanding such considerations, it is worthwhile to point out that the retained potential morbidity-related biomarkers, as determined from this analysis are suggestive that the magnitude of inflammatory responses to OSA is a major contributing mechanisms to end-organ morbidity. Thus, these findings further reinforce the now well-established conceptual framework that OSA is a low grade systemic inflammatory disease.

In summary, this scoping review provides insights into our current state of knowledge about biological markers and their applicability in the detection of OSA-induced morbidity. Over the last 12 years, a substantial number of studies have aimed to identify an ideal biomarker for OSA morbidity. The most promising biomarkers identified revolved around the identification of CVD in OSA patients. The 48 studies identified in this scoping review should be helpful in identifying appropriate questions in the field and formulate adequate research strategies in this important area.

Limitations

The absence of a unique PSG criterion for the diagnosis OSA may lead to different interpretations of the findings in the studies included in this review. The studies included here explored different types of morbidities such that the critical mass of studies addressing a specific morbidity is unlikely. Most studies used a clinical referral sample with no *post-hoc* verification in

more generalized community settings. Importantly, the duration of OSA was now evaluated and included as a potential confounder to the validity of any of the putative biomarkers. Furthermore, considering that some gender-related differences in urinary biomarkers aiming at diagnosing OSA in children have been reported,⁷¹ it would be interesting and potentially relevant to explore gender as a modifier of the receiver-operator curve properties of selected biomarkers. Finally, the majority of studies have been performed in the United States, such that extrapolation to other populations is uncertain.

5 Conclusions

Blood-based biomarkers accounted for the majority of the studies, and most of the explored approaches did not identify definitive biomarkers of OSA morbidity. IL-6 and hsCRP appear to exhibit a favorable profile as biomarkers aiming to discriminate OSA patients with and without morbidity in adults, as well as MRP 8/14 in children. Urinary neurotransmitters can potential be a good tool for screening cognitive function in OSA children.

References

1. Leger D, Bayon V, Laaban JP, Philip P. Impact of sleep apnea on economics. *Sleep medicine reviews*. Oct 2012;16(5):455-462.
2. Kim J, Bhattacharjee R, Kheirandish-Gozal L, Spruyt K, Gozal D. Circulating microparticles in children with sleep disordered breathing. *Chest*. Aug 2011;140(2):408-417.
3. Del Ben M, Fabiani M, Loffredo L, et al. Oxidative stress mediated arterial dysfunction in patients with obstructive sleep apnoea and the effect of continuous positive airway pressure treatment. *BMC pulmonary medicine*. 2012;12:36.
4. Bostrom KB, Hedner J, Melander O, et al. Interaction between the angiotensin-converting enzyme gene insertion/deletion polymorphism and obstructive sleep apnoea as a mechanism for hypertension. *Journal of hypertension*. Apr 2007;25(4):779-783.
5. Montesi SB, Bajwa EK, Malhotra A. Biomarkers of sleep apnea. *Chest*. Jul 2012;142(1):239-245.
6. Kanbay A, Kaya E, Buyukoglan H, et al. Serum gamma-glutamyl transferase activity is an independent predictor for cardiovascular disease in obstructive sleep apnea syndrome. *Respiratory medicine*. Apr 2011;105(4):637-642.
7. Celec P, Hodosy J, Behuliak M, et al. Oxidative and carbonyl stress in patients with obstructive sleep apnea treated with continuous positive airway pressure. *Sleep & breathing = Schlaf & Atmung*. Jun 2012;16(2):393-398.
8. Gozal D, Crabtree VM, Sans Capdevila O, Witcher LA, Kheirandish-Gozal L. C-reactive protein, obstructive sleep apnea, and cognitive dysfunction in school-aged children. *American journal of respiratory and critical care medicine*. Jul 15 2007;176(2):188-193.
9. Kelly A, Dougherty S, Cucchiara A, Marcus CL, Brooks LJ. Catecholamines, adiponectin, and insulin resistance as measured by HOMA in children with obstructive sleep apnea. *Sleep*. Sep 2010;33(9):1185-1191.
10. Papanas N, Steiropoulos P, Nena E, et al. HbA1c is associated with severity of obstructive sleep apnea hypopnea syndrome in nondiabetic men. *Vascular health and risk management*. 2009;5:751-756.

11. Drager LF, Togeiro SM, Polotsky VY, Lorenzi-Filho G. Obstructive sleep apnea: a cardiometabolic risk in obesity and the metabolic syndrome. *Journal of the American College of Cardiology*. Aug 13 2013;62(7):569-576.
12. Pastore AL, Palleschi G, Ripoli A, et al. Severe obstructive sleep apnoea syndrome and erectile dysfunction: a prospective randomised study to compare sildenafil vs. nasal continuous positive airway pressure. *International journal of clinical practice*. Aug 2014;68(8):995-1000.
13. Sans Capdevila O, Crabtree VM, Kheirandish-Gozal L, Gozal D. Increased morning brain natriuretic peptide levels in children with nocturnal enuresis and sleep-disordered breathing: a community-based study. *Pediatrics*. May 2008;121(5):e1208-1214.
14. Gozal D, Serpero LD, Kheirandish-Gozal L, Capdevila OS, Khalyfa A, Tauman R. Sleep measures and morning plasma TNF-alpha levels in children with sleep-disordered breathing. *Sleep*. Mar 2010;33(3):319-325.
15. Sanchez-de-la-Torre M, Barcelo A, Pierola J, et al. Plasma levels of neuropeptides and metabolic hormones, and sleepiness in obstructive sleep apnea. *Respiratory medicine*. Dec 2011;105(12):1954-1960.
16. Tarasiuk A, Reuveni H. The economic impact of obstructive sleep apnea. *Current opinion in pulmonary medicine*. Nov 2013;19(6):639-644.
17. Baldi I, Gulati A, Lorenzoni G, et al. Public health implications of obstructive sleep apnea burden. *Indian journal of pediatrics*. Sep 2014;81 Suppl 1:55-62.
18. Kuhle S, Kirk S, Ohinmaa A, Yasui Y, Allen AC, Veugelers PJ. Use and cost of health services among overweight and obese Canadian children. *International journal of pediatric obesity : IJPO : an official journal of the International Association for the Study of Obesity*. Apr 2011;6(2):142-148.
19. Yamagishi K, Ohira T, Nakano H, et al. Cross-cultural comparison of the sleep-disordered breathing prevalence among Americans and Japanese. *The European respiratory journal*. Aug 2010;36(2):379-384.
20. Lumeng JC, Chervin RD. Epidemiology of pediatric obstructive sleep apnea. *Proceedings of the American Thoracic Society*. Feb 15 2008;5(2):242-252.
21. EO B. Sleep disordered breathing in children in a general population sample: prevalence and risk factors. *Sleep*. 2009;32(6):731-736.

22. Kheirandish-Gozal L. What is "abnormal" in pediatric sleep? *Respiratory care*. Oct 2010;55(10):1366-1374; discussion 1374-1366.
23. Shih JL, Malhotra A. Could vitamins be helpful to patients with sleep apnea? *Chest*. Feb 2011;139(2):237-238.
24. Arksey H OM, L. Scoping studies: towards a methodological framework. *International Journal of Social Research Methodology*. 2005;8(1):19-32.
25. Marcus CL, Brooks LJ, Draper KA, et al. Diagnosis and management of childhood obstructive sleep apnea syndrome. *Pediatrics*. Sep 2012;130(3):576-584.
26. El-Solh AA, Mador MJ, Sikka P, Dhillon RS, Amsterdam D, Grant BJ. Adhesion molecules in patients with coronary artery disease and moderate-to-severe obstructive sleep apnea. *Chest*. May 2002;121(5):1541-1547.
27. Jordan W, Hagedohm J, Wiltfang J, et al. Biochemical markers of cerebrovascular injury in sleep apnoea syndrome. *European Respiratory Journal*. 2002;20(1):158-164.
28. Gami AS, Svatikova A, Wolk R, et al. Cardiac troponin T in obstructive sleep apnea. *Chest*. Jun 2004;125(6):2097-2100.
29. Kokturk O, Ciftci TU, Mollarecep E, Ciftci B. Elevated C-reactive protein levels and increased cardiovascular risk in patients with obstructive sleep apnea syndrome. *International heart journal*. Sep 2005;46(5):801-809.
30. Kokturk O, Ciftci TU, Mollarecep E, Ciftci B. Serum homocysteine levels and cardiovascular morbidity in obstructive sleep apnea syndrome. *Respiratory medicine*. Mar 2006;100(3):536-541.
31. Saletu M, Nosiska D, Kapfhammer G, et al. Structural and serum surrogate markers of cerebrovascular disease in obstructive sleep apnea (OSA): association of mild OSA with early atherosclerosis. *Journal of neurology*. Jun 2006;253(6):746-752.
32. El Solh AA, Akinnusi ME, Baddoura FH, Mankowski CR. Endothelial cell apoptosis in obstructive sleep apnea: a link to endothelial dysfunction. *American journal of respiratory and critical care medicine*. Jun 1 2007;175(11):1186-1191.
33. Minoguchi K, Yokoe T, Tazaki T, et al. Silent brain infarction and platelet activation in obstructive sleep apnea. *American journal of respiratory and critical care medicine*. Mar 15 2007;175(6):612-617.

34. Bravo Mde L, Serpero LD, Barcelo A, Barbe F, Agusti A, Gozal D. Inflammatory proteins in patients with obstructive sleep apnea with and without daytime sleepiness. *Sleep & breathing = Schlaf & Atmung*. Sep 2007;11(3):177-185.
35. da Silva LG, Mottin CC, Souza DO, et al. Serum S100B but not NSE levels are increased in morbidly obese individuals affected by obstructive sleep apnea-hypopnea syndrome. *Obesity surgery*. Aug 2008;18(8):993-999.
36. Hubner RH, El Mokhtari NE, Freitag S, et al. NT-proBNP is not elevated in patients with obstructive sleep apnoea. *Respiratory medicine*. Jan 2008;102(1):134-142.
37. Mills PJ, Kim JH, Bardwell W, Hong S, Dimsdale JE. Predictors of fatigue in obstructive sleep apnea. *Sleep & breathing = Schlaf & Atmung*. Nov 2008;12(4):397-399.
38. Ursavas A, Karadag M, Oral AY, Demirdogen E, Oral HB, Ege E. Association between serum neopterin, obesity and daytime sleepiness in patients with obstructive sleep apnea. *Respiratory medicine*. Aug 2008;102(8):1193-1197.
39. Al Lawati N, Mulgrew A, Cheema R, et al. Pro-atherogenic cytokine profile of patients with suspected obstructive sleep apnea. *Sleep & breathing = Schlaf & Atmung*. Nov 2009;13(4):391-395.
40. Koyama RG, Drager LF, Lorenzi-Filho G, et al. Reciprocal interactions of obstructive sleep apnea and hypertension associated with ACE I/D polymorphism in males. *Sleep medicine*. Dec 2009;10(10):1107-1111.
41. Culla B, Guida G, Brussino L, et al. Increased oral nitric oxide in obstructive sleep apnoea. *Respiratory medicine*. Feb 2010;104(2):316-320.
42. Aihara K, Oga T, Harada Y, et al. Comparison of biomarkers of subclinical lung injury in obstructive sleep apnea. *Respiratory medicine*. Jun 2011;105(6):939-945.
43. Bekci TT, Kayrak M, Kiyici A, et al. The association among lipoprotein-associated phospholipase A2 levels, total antioxidant capacity and arousal in male patients with OSA. *International journal of medical sciences*. 2011;8(5):369-376.
44. Kato K, Takata Y, Usui Y, et al. Severe obstructive sleep apnea increases cystatin C in clinically latent renal dysfunction. *Respiratory medicine*. Apr 2011;105(4):643-649.
45. Mohsenin V, Urbano F. Circulating antiangiogenic proteins in obstructive sleep apnea and hypertension. *Respiratory medicine*. May 2011;105(5):801-807.

46. Rahangdale S, Yeh SY, Novack V, et al. The influence of intermittent hypoxemia on platelet activation in obese patients with obstructive sleep apnea. *Journal of clinical sleep medicine : JCSM : official publication of the American Academy of Sleep Medicine*. Apr 15 2011;7(2):172-178.
47. Zhao Q, Liu ZH, Zhao ZH, et al. Effects of obstructive sleep apnea and its treatment on cardiovascular risk in CAD patients. *Respiratory medicine*. Oct 2011;105(10):1557-1564.
48. Cherneva RV, Georgiev OB, Petrova DS, et al. Resistin - the link between adipose tissue dysfunction and insulin resistance in patients with obstructive sleep apnea. *Journal of diabetes and metabolic disorders*. 2013;12(1):5.
49. Cholidou KG, Kostakis ID, Manali ED, et al. Calprotectin: a protein related to cardiovascular risk in adult patients with obstructive sleep apnea. *Cytokine*. Mar 2013;61(3):917-923.
50. Sales LV, Bruin VM, D'Almeida V, et al. Cognition and biomarkers of oxidative stress in obstructive sleep apnea. *Clinics*. 2013;68(4):449-455.
51. Salord N, Gasa M, Mayos M, et al. Impact of OSA on biological markers in morbid obesity and metabolic syndrome. *Journal of clinical sleep medicine : JCSM : official publication of the American Academy of Sleep Medicine*. Mar 15 2014;10(3):263-270.
52. Oyama J, Yamamoto H, Maeda T, Ito A, Node K, Makino N. Continuous positive airway pressure therapy improves vascular dysfunction and decreases oxidative stress in patients with the metabolic syndrome and obstructive sleep apnea syndrome. *Clinical cardiology*. Apr 2012;35(4):231-236.
53. Paik MJ, Kim DK, Nguyen DT, et al. Correlation of daytime sleepiness with urine metabolites in patients with obstructive sleep apnea. *Sleep & breathing = Schlaf & Atmung*. Sep 2014;18(3):517-523.
54. Biltagi MA, Maguid MA, Ghafar MA, Farid E. Correlation of 8-isoprostane, interleukin-6 and cardiac functions with clinical score in childhood obstructive sleep apnoea. *Acta paediatrica*. Oct 2008;97(10):1397-1405.
55. Wang Y, Wang JJ, Zhao MQ, Liu SM, Li YZ. Changes of Serum Brain-Derived Neurotrophic Factor in Children with Obstructive Sleep Apnoea-Hypopnoea Syndrome following Adenotonsillectomy. *Journal of International Medical Research*. 2010;38(6):1942-1951.
56. Gozal D, Kheirandish-Gozal L, Bhattacharjee R, Spruyt K. Neurocognitive and endothelial dysfunction in children with obstructive sleep apnea. *Pediatrics*. Nov 2010;126(5):e1161-1167.

57. Kim J, Lee S, Bhattacharjee R, Khalyfa A, Kheirandish-Gozal L, Gozal D. Leukocyte telomere length and plasma catestatin and myeloid-related protein 8/14 concentrations in children with obstructive sleep apnea. *Chest*. Jul 2010;138(1):91-99.
58. Kheirandish-Gozal L, Bhattacharjee R, Kim J, Clair HB, Gozal D. Endothelial progenitor cells and vascular dysfunction in children with obstructive sleep apnea. *American journal of respiratory and critical care medicine*. Jul 1 2010;182(1):92-97.
59. Kheirandish-Gozal L, McManus CJ, Kellermann GH, Samiei A, Gozal D. Urinary neurotransmitters are selectively altered in children with obstructive sleep apnea and predict cognitive morbidity. *Chest*. Jun 2013;143(6):1576-1583.
60. Kim J, Bhattacharjee R, Snow AB, Capdevila OS, Kheirandish-Gozal L, Gozal D. Myeloid-related protein 8/14 levels in children with obstructive sleep apnoea. *The European respiratory journal*. Apr 2010;35(4):843-850.
61. Gozal D, Sans Capdevila O, McLaughlin Crabtree V, Serpero LD, Witcher LA, Kheirandish-Gozal L. Plasma IGF-1 levels and cognitive dysfunction in children with obstructive sleep apnea. *Sleep medicine*. Feb 2009;10(2):167-173.
62. Gozal D, Capdevila OS, Kheirandish-Gozal L, Crabtree VM. APOE epsilon 4 allele, cognitive dysfunction, and obstructive sleep apnea in children. *Neurology*. Jul 17 2007;69(3):243-249.
63. Papanas N, Steiropoulos P, Nena E, et al. Predictors of obstructive sleep apnea in males with metabolic syndrome. *Vascular health and risk management*. 2010;6:281-286.
64. Jullian-Desayes I, Joyeux-Faure M, Tamisier R, et al. Impact of obstructive sleep apnea treatment by continuous positive airway pressure on cardiometabolic biomarkers: A systematic review from sham CPAP randomized controlled trials. *Sleep medicine reviews*. Jul 31 2014.
65. Gozal D. Serum, urine, and breath-related biomarkers in the diagnosis of obstructive sleep apnea in children: is it for real? *Current opinion in pulmonary medicine*. 2012;18:561-567.
66. Wong TK. The search on an ideal disease marker for childhood obstructive sleep apnea syndrome. *Sleep*. Feb 2011;34(2):133-134.
67. Gozal D, Kheirandish-Gozal L. New approaches to the diagnosis of sleep-disordered breathing in children. *Sleep medicine*. Aug 2010;11(7):708-713.

68. International classification of sleep disorders. Diagnostic and coding manual *Section on sleep related bruxism*. Illinois: American Academy of Sleep Medicine; 2005:189-192.
69. Qaseem A, Dallas P, Owens DK, et al. Diagnosis of obstructive sleep apnea in adults: a clinical practice guideline from the american college of physicians. *Annals of internal medicine*. Aug 5 2014;161(3):210-220.
70. Brockmann PE, Schaefer C, Poets A, Poets CF, Urschitz MS. Diagnosis of obstructive sleep apnea in children: a systematic review. *Sleep medicine reviews*. Oct 2013;17(5):331-340.
71. Becker L, Kheirandish-Gozal L, Peris E, Schoenfelt KQ, Gozal D. Contextualized urinary biomarker analysis facilitates diagnosis of pediatric obstructive sleep apnea. *Sleep Medicine* 2014; 15:541-549.

Figure 1 - Flow Diagram of literature search and selection criteria¹.

¹ [Adapted from PRISMA.](#)

Table 1 - Summary of study descriptive characteristics of included pediatric articles (n=6). The biomarker clinical application was classified as (1) potential morbidity biomarker(s) in OSA patients, (2) inconclusive as biomarker, and (3) evidence not supportive as potential biomarker(s). The level of evidence was classified in A (well designed prognostic or diagnostic studies on relevant population), B (prognostic or diagnostic studies with minor limitations, overwhelmingly consistent evidence from observational studies), C (observational studies (case-control and cohort design) (n=14)

Year	Author	Country	Cases	Control	Mean Age or range if provided	Morbidity	Biomarker Type	OSA diagnostic criteria (PSG)
2007	Gozal et al ⁶²	United States	Snoring children divided in 2 groups: Non-OSA (n=112/62 male) and OSA (n=146/79 male)	Non-snoring children (n=87/47 male)	5-7	Cognitive dysfunction	Blood	OAI > 1 and AHI > 2
2007	Gozal et al ⁸	United States	Total (n=205) Snoring Non-OSA (n=103/57 male) Snoring OSA (n=102/59 male)	Non-snoring with AHI < 1 (n=73/40 male)	5-7	Cognitive dysfunction	Blood	AHI > 1 and/or AHI > 2
2008	Biltagi et al ⁵⁴	Egypt	Children with adenoidal hypertrophy OSA suspected (n=40)	Healthy children (n=20)	4-8	Cardiac Diastolic dysfunction	EBC	clinical assessment score ≥ 40
2008	Sans Capdevila et al ¹³	United States	OSA children (n=20) HS children (n=20)	Controls (n=20)	5-7	Enuresis		OAI > 1 and AHI > 2
2009	Gozal et al ⁶¹	United States	Snoring children: Non-OSA (n=23/13 male) OSA (n=87/46 male)	Non-snoring (n=52/28 male)	5-7	Cognitive dysfunction	Blood	OAI > 1 and/or AHI > 2 with a nadir oxygen saturation value at least < 92%
2010	Gozal et al ¹⁴	United States	OSA children (n=15)	Non-OSA (n=15)	7.3#	EDS	Blood	AHI ≥ 1
2010	Gozal et al ⁵⁶	United States	OSA children (n=87/51 male)	Non-snoring (n=21/11 male)	5-8	Neurocognitive and Endothelial Dysfunction	Blood	OAI ≥ 1 and AHI ≥ 5
2010	Kelly et al ⁹	United States	OSA suspected children: Pre-pubertal (n=20/11 male) Pubertal (n=34/21 male)	Non-OSA Pre-pubertal (n=17/12 male) Pubertal (n=27/12 male)	4-18	Insulin Resistance	Blood /urine	AHI > 1.5
2010	Kheirandish-Gozal et al ⁵⁸	United States	OSA children (n=80)	Non-OSA (n=20)	8.2	Vascular Dysfunction	Blood	OAI ≥ 1 and AHI ≥ 5
2010	Kim et	United	OSA (n=140/85 male#)	Non-OSA	7.6	Endothelial	Blood	AHI ≥ 1

	al ⁶⁰	States	(n=115/64 male#)			dysfunction		
2010	Kim et al ⁵⁷	United States	Mild OSA (n=85) Moderate-to-severe OSA (n=26)	Non-OSA (n=102)	7.7	emtabolic and endothelial dysfunction	Blood	AHI \geq 1
2010	Wang et al ⁵⁵	China	OSA (n=44/28 male)	Non-OSA (n=26/15 male)	6.3#	Cognitive dysfunction	Blood	RDI \geq 5
2011	Kim et al ²	United States	Mild OSA (n=65/42 male) Moderate to severe OSA (n=14/9 male)	Non-OSA (n=56/30 male)	4-12	Endothelial dysfunction	Blood	AHI $>$ 1
2013	Kheirandish-Gozal et al ⁵⁹	United States	OSA children (n=50/27 male#))	Non-OSA (n=20/11 male#)	6-12	Cognitive dysfunction	Urine	AHI \geq 2

All terms that mean obstructive sleep apnea (SDB, SRDB, OSAS) were standardized as OSA. # Mean calculated by author.

A&T=adenotonsillectomy, AHI=apnea/hypopnea index, APO=apolipoprotein, BMI=body mass index, BNP=brain natriuretic peptide levels, BDNF= Brain-Derived Neurotrophic Factor, cICAM-1=circulating intercellular adhesion molecule 1, CRP=C reactive protein, EBC= exhaled breath condensate, EDS=excessive daytime sleepiness, EPCs=endothelial progenitor cells, Hs-CRP= high sensitivity C-reactive protein, IGF=insulin growth factor, IL-6=interleukin-6, MPs=microparticles, MRP=myeloid-related protein, OSA=obstructive sleep apnea, RDI= respiratory disturbance index, TNF- α = tumor necrosis factor alpha.

Table 2 - Summary of study descriptive characteristics of included articles (adults, n=33). The biomarker clinical application was classified as (1) potential biomarker(s) of OSA-associated morbidity; (2) inconclusive for morbidity biomarker, and (3) evidence not supportive as potential morbidity biomarker(s). The level of evidence was classified in A (well designed prognostic or diagnostic studies on relevant population), B (prognostic or diagnostic studies with minor limitations, overwhelmingly consistent evidence from observational studies), C (observational studies (case-control and cohort design) (n=34)

Year	Author	Country	Cases	Control	Mean Age or range if provided	Morbidity	Type of Biomarker	OSA diagnostic criteria (PSG)
2002	El-Solh et al ²⁶	United States	Patients with moderate to severe OSA (n=15) matched with controls by age, gender, BMI, and severity of CAD	Non-OSA (n=15)	61.2	CAD	Blood	AHI $\geq 10/h$
2002	Jordan et al ²⁷	Germany	OSA patients (n=19/19 male) divided in 2 groups: Group 1: patients with OSA without severe hypoxemia during sleep Group 2: patients with OSA with severe hypoxemia during sleep	-	50.6	Cerebrovascular injury	Blood	RDI > 40/h
2004	Gami et al ²⁸	United States	Patients with moderate to severe OSA (n=15/15 male)	-	62.0	CAD	Blood	AHI > 15/h
2005	Kokturk et al ²⁹	Turkey	Group 1: Patients with OSA and CVD (n=38) Group 2: Patients with OSA without CVD (n=56)	Group 3: patients without OSA with CVD (n=57)	45.0#	Cardiovascular morbidity	Blood	AHI $\geq 5/h$
2006	Kokturk et al ³⁰	Turkey	Group 1: OSA obese patients with CVD (n=25) Group 2: OSA patients without CVD (n=47) Matched with controls by age and BMI	Group 3: Non-OSA obese with CVD (n=42)	49.7#	Cardiovascular morbidity	Blood	AHI $\geq 5/h$
2006	Saletu et al ³¹	Austria	Group 1: mild OSA (n=27) Group 2: moderate OSA (n=25) Group 3: severe OSA (n=51)	Non-OSA (n=44)	53.5#	Cerebrovascular disease	Blood	AHI $\geq 5/h$
2007	Bostrom et al ⁴	Sweden	Hypertensive patients in primary care (n=157)	Normotensive	61.0#	Hypertension and cardiovascular	Blood	AHI $\geq 10/h$

				populatio n controls (n=181)		complications		
2007	El Solh et al ³²	United States	OSA (n=14)	Non- OSA (n=10)	44.4#	Atherosclerosis	Blood	AHI>5/h
2007	Minoguc hi et al ³³	Japan	OSA (n=50/50 male) Matched with control by age and weight	Non- OSA obese patients (n=15)	48.9#	Cerebrovascular disease	Blood	AHI≥5/h
2007	Pena Bravo et al ³⁴	Spain	OSA (n=50/50 male) Matched by age with controls Without personal or family history of CVD or diabetes	Non- OSA (n=20)	50.3#	Daytime sleepiness	Blood	AHI≥20/ h
2008	Da Silva et al ³⁵	Brazil	Morbidity obese (n=25/10 male) with OSA complaints	-	39.9	Cerebral injury	Blood	AHI≥5/h
2008	Hubner et al ³⁶	Germany	OSA suspected (n=60)	-	55.7	Cardiovascular stress and dysfunction	Blood	AHI≥5/h
2008	Mills et al ³⁷	South Korea	Untreated OSA (n=56/41 male)	-	49.2	Fatigue	Blood	AHI≥15/ h
2008	Ursavas et al ³⁸	Turkey	OSA (n=22)	Non- OSA (n=18)	47.8#	Obesity	Blood	AHI≥5/h
2009	Al Lawati et al ³⁹	Canada	OSA (n=176/119 male)	-	50.0	Cardiovascular disease	Blood	AHI≥5/h
2009	Koyama et al ⁴⁰	Brazil	OSA (n=266/266 male) divided in 2 groups: Normotensive (n=152) Hypertensive (n=114)	-	48.0	Angiotensin- converting enzyme (ACE) polymorphism	Blood	AHI>5/h
2009	Papanas et al ¹¹	Greece	OSA (n=31/31 male)	-	46.7	Diabetes	Blood	AHI>15/ h
2010	Culla et al ⁴¹	Italy	OSA (n=39)	Non- OSA non smoking with	53.8#	Oropharyngeal inflammation	EBC	AHI>5/h

				CRS (n=15) and Non- OSA non- smoking with mild asthma (n=26)				
2011	Aihara et al ⁴²	Japan	OSA: Mild OSA (n=25) Moderate OSA (n=52) Severe OSA (n=73)	Non- OSA (n=20)	53.8#	Lung injury	Blood	AHI>5/h
2011	Bekci et al ⁴³	Turkey	Group 1: Below the median arousal Group 2: Above the median arousal	-	43.5	Cardiac and vascular events	Blood	AHI≥5/h
2011	Kanbay et al ⁶	Turkey	OSA Mild OSA (n=17) Moderate OSA (n=34) Severe OSA (n=93)	Non- OSA (n=22)	54.2#	Cardiovascular disease	Blood	AHI>5/h
2011	Kato et al ⁴⁴	Japan	(n=267/233 male)	-	51.0	Renal dysfunction	Blood	AHI≥5/h
2011	Mohseni n and Urbano ⁴⁵	United States	OSA Normotensive OSA (n=11/10 male) Hypertensive OSA (n=11/10 male)	-	46.5#	Hypertension	Blood	AHI≥30/ h
2011	Rahangdale et al ⁴⁶	United States	OSA obese non-smokers (n=47) Matched with controls by BMI	Non- OSA obese non- smokers (n=30)	38.7#	Platelet activation	Blood	AHI≥10/ h
2011	Sanchez-de-la torre et al ¹⁵	Spain	OSA with EDS (n=132/110 male) Matched by age, gender, BMI, AHI with controls	OSA without EDS (n=132/113 male)	50.5#	EDS	Blood	AHI≥20/ h
2011	Zhao et al ⁴⁷	Australia	OSA Mild OSA (n=50/45 male) Moderate OSA (n=43/38 male)	Non- OSA (n=25/20 male)	58.7#	Coronary artery disease	Blood	AHI≥5/h

			Severe OSA (n=33/27 male)					
2012	Celec et al ⁷	Slovakia	OSA (n=89/67 male)	-	56.1	Cardiovascular disease	Blood, saliva	RDI>30
2012	Del Ben et al ³	Italy	OSA Mild/moderate OSA (n=61) Severe OSA (n=30)	Non-OSA snorers (n=47)	53.7#	Endothelial dysfunction and oxidative stress	Blood, urine	(home PSG) AHI≥5/h
2012	Oyama et al ⁵²	Japan	OSA + Mets patients (n=32/19 male) who received CPAP therapy for 3 months	-	53.9	Metabolic syndrome	Blood	AHI>20/h
2013	Cherneva et al ⁴⁸	Bulgaria	OSA: Diabetic OSA patients (n=23/17 male) Impaired glucose tolerance (n=27/23 male) Normal glucose metabolism (n=17/17 male)	-	49.8#	Diabetes	Blood, urine	AHI≥5/h
2013	Cholidou et al ⁴⁹	Greece	OSA Moderate OSA (n=33/25 male) Severe OSA (n=27/26 male)	Non-OSA (n=14/12 male)	45.2#	Cardiovascular disease	Blood	AHI>5/h
2013	Sales et al ⁵⁰	Brazil	OSA (n=14) Matched by years of schooling, age, BMI with controls	Non-OSA (n=13)	36.0	Neuropsychological performance and oxidative stress	Blood	AHI>10/h and at least one symptom or AHI >15/h
2014	Paik et al ⁵³	South Korea	Sleepy OSA (n=26/19 male)	Non-sleepy OSA (n=23/18 male)	45.5#	EDS	Urine	AHI>5
2014	Salord et al ⁵¹	Spain	Mets Mets Non-OSA (n=13/5 male) Mets OSA (n=26/6 male)	Non-OSA Non-Mets (n=13/4 male)	43.0#	Mets	Blood	AHI≥15/h

All terms that mean obstructive sleep apnea (SDB, SRDB, OSAS) were standardized as OSA. # Mean calculated by author.

8-iso-PGF₂α=8-iso-prostaglandin F₂ α, ACE=Angiotensin converting enzyme, ADMA=asymmetrical dimethylarginine, AGEs=advanced glycation end products, AHI=apnea/hypopnea index, AOPP=advanced oxidation protein products, BMI=body mass index, CAD=coronary artery disease, CD146=melanoma cell adhesion molecule, CPAP=continuous positive airway pressure, CRP=C reactive protein, CRS=chronic rhinitis or rhinosinusitis,

CVD=cardiovascular disease, EBC=exhaled breath condensate, EDS=excessive daytime sleepiness, ET-1=endothelin-1, FENO=exhaled nitric oxide, FFA=free fatty acids, FRAP= ferric reducing antioxidant power, FRAS= ferric-reducing ability of serum, GGT=serum gamma-glutamyl transferase, GPIIb/IIIa= glycoprotein IIb/IIIa, HbA_{1c}=glycosylated hemoglobin, HDL=high density lipoprotein, Hs-CRP=high sensitivity C-reactive protein, I/D=insertion/deletion, ICAM-1=intercellular adhesion molecule 1, IL-1B=interleukin-1B, IL-6=interleukin-6, IL-8=interleukin-8, KL-6=Krebs Von den Lungen-6, LDL= low density lipoprotein, Lp-PLA2=lipoprotein-associated phospholipase A2, Mets=morbid obesity and metabolic syndrome, NO_x=Nitrogen oxide, NP=neopterin, NSE=neuron-specific enolase (a glycolytic pathway enzyme from CNS), NT-ProBNP=N-terminal pro-brain natriuretic peptide, oNO=oral nitric oxide, OSA=obstructive sleep apnea, P-selectin= Platelet selectin, PGF2- α =Prostaglandin F2 α , PSG=polysomnography, RDI=respiratory disturbance index, S100B=calcium binding protein B, SBI=silent brain infarction, sCD40= soluble CD40, sCD40L=soluble CD40 ligand, sEng=soluble endoglin, sFasL= soluble Fas ligand, sFlt-1=soluble fms-like tyrosine kinase-1, sICAM-1=soluble Inercellular adhesion molecule 1, sNOX2-dp=soluble NOX2 derived peptide, SP-D=surfactant protein, sP-selectin=soluble P-selectin, sTNF-RI=soluble tumor necrosis factor receptor 1, sVCAM-1= soluble vascular cell adhesion molecule-1, TAC=total antioxidant capacity , TBARS=thiobarbituric acid reactive substance, TNF- α =tumor necrose factor alpha, VCAM-1=vascular cell adhesion molecule-1, VEGF=vascular endothelial growth factor.

Table 3 - Potential biomarkers identified (n=31)

Potential Biomarkers	Children
8-isoprostane	✓
Adiponectin	✓
AGEs	
AOPP	
APOEε4	✓
Calprotectin	
Catecholamines	✓
Catestatin	✓
CRP	✓
Cystatin C	
FFA	
FRAP	
FRAS	
Fructosamine	
GGT	
IL-6	✓
IL-8	
HOMA	✓
KL-6	
Leptin	
Lipid profiles	
MRP8/14	✓
NP	
Resistin	
sTNF-RI	
S100B	
sVCAM-1	
TAC	
TBARS	
TNF- α	✓
Urinary Neurotransmitters	✓

AGEs=advanced glycation end products, AOPP=advanced oxidation protein products, hsCRP=high sensitivity C-reactive protein, FFA= free fatty acids, FRAP=ferric reducing antioxidant power, FRAS=ferric reducing ability of serum, GGT=serum gamma-glutamyl transferase, IL-1B=interleukin 1B, IL-6=interleukin-6, IL-8=interleukin-8, HOMA=homeostatic model assessment of insulin resistance, KL-6= Krebs von den Lungen-6, MRP=myeloid-related protein, NP=neopterin, S100B=calcium binding protein B, sVCAM-1=soluble vascular cell adhesion molecule-1, TAC=total antioxidant capacity, TBARS=thiobarbituric acid reactive substance, TNF-α=tumor necrosis factor alpha.

Table 4 - Distribution of studies according to country (n=17).

Country	Total	Children	Adults
United States	17	12	5
Turkey	5	0	5
Japan	4	0	4
Brazil	3	0	3
Spain	3	0	3
Germany	2	0	2
Greece	2	0	2
Italy	2	0	2
Australia	1	0	1
Austria	1	0	1
Bulgaria	1	0	1
Canada	1	0	1
China	1	1	0
Egypt	1	1	0
Slovakia	1	0	1
South Korea	2	0	2
Sweden	1	0	1
Total	48	14	34